

PolyMem®

CASE STUDY

Persistent Multiple Venous Dermatitis Ulcerations Closed Completely Using PolyMem Wic Silver® and PolyMem Dressings Under Compression



BEFORE



AFTER



SIX MONTHS POST DISMISSAL

Persistent Multiple Venous Dermatitis Ulcerations Closed Completely Using PolyMem Wic Silver and PolyMem Dressings Under Compression

Janis Harrison, RN, BSN, CWOCN, CFCN, Pender Community Hospital, 603 Earl Street, Pender, NE 68047

PROBLEM

A 69-year-old female with venous hypertension dermatitis ulcerations had a history of arthritis treated with regular aspirin, COPD treated with albuterol, hypertension treated with a diuretic, varicose vein ligation and recent bilateral knee surgery. She presented with pneumonia and several small pruritic open areas on both ankles. The patient had self-treated these recurring wounds with a steroid ointment. Then they were treated by her doctor with silver sulfadiazine covered with 4 x 4's and wrapped with absorbent stretch gauze for a month. The ankle areas were warm to the touch and draining.

The WOC Nurse was consulted when the patient was admitted to the hospital with severe itching, pain, and cellulitis of the right lower leg. ABI's: Left 0.86 mm Hg; Right 0.80 mm Hg. Hospital treatment included daily whirlpool cleansing, barrier cream to the periwound skin, and dressing all open areas with PolyMem Wic Silver cavity filler covered with absorbent stretch gauze and then two layers of compression stocking material. The patient kept the leg elevated in bed. After a 10 day hospital stay, an Unna's Boot was applied by the physician and the patient was dismissed to the out-patient wound clinic, where she was seen 1 week later by the WOC Nurse. At this point the PolyMem Wic cavity filler and 4-layer compression wrap treatment was initiated.

RATIONALE

Heavily draining venous ulcer wounds treated with PolyMem Wic cavity filler under compression closed very quickly compared with wounds treated without this cavity filler in previous patients. Excess exudate is wicked directly away from the wound surface, through the PolyMem Wic cavity filler into the absorptive layers of the compression dressing, thus increasing wear time while preventing maceration. The PolyMem Wic cavity filler contains glycerin to soothe and hydrate the wound while drawing and concentrating healing substances from the body to the wound bed. It also contains a gentle surfactant to continually cleanse the wound bed. The delicate new structures in the wound bed are preserved because no additional cleansing is needed during dressing changes. This treatment regimen seemed to be an ideal choice for this patient.

METHODOLOGY

Initially the wound and surrounding area were cleansed and PolyMem Wic cavity filler covered with a 4-layer compression wrap was applied, to be changed every 5 – 7 days. The wound bed was not routinely cleansed with subsequent dressing changes. After 6 weeks the right lateral wounds closed and the remaining wounds were very small with scant exudate, so the wounds were covered with PolyMem Wic dressings and the patient was fitted with 20 – 30 mm Hg compression stockings.

A month later, new ulcers formed, which were MRSA positive. Oral antibiotic therapy was initiated. The treatment protocol of PolyMem Wic cavity filler covered with a 4-layer compression wrap was resumed, with barrier cream to periwound areas. Three weeks of this treatment resulted in almost complete wound closure again, but due to the chronicity of the wounds, this time the treatment was continued until complete closure was achieved. The patient was measured for thigh-high compression stockings 20 – 30 mm Hg.

Three days later the patient was again hospitalized, this time for IV antibiotic therapy due to a MRSA infection on her hand (injured from scratching at her dressing) and her right lower leg. The treatment was modified to include PolyMem Wic Silver instead of PolyMem Wic cavity filler, covered with the 4-layer compression dressing. Five days after this change, the wounds were healing and most were closed. PolyMem Wic cavity filler was again applied and covered with a 4-layer compression wrap. This was continued 10 more days, until all areas were closed. At that point the patient was placed on long term oral antibiotics for 3 months and dismissed from wound care. The patient applied skin conditioning cream and wore thigh-high compression stockings 20 – 30 mm Hg.

RESULTS

The protocol of PolyMem Wic cavity filler applied on the wound and covered with a 4-layer compression wrap was successful in closing these venous dermatitis induced wounds in 3 - 6 weeks. But, the MRSA infection relapses greatly prolonged the patient's overall recovery time. Initially the patient rated her pain a "10" on the 0-10 scale. By the second dressing change the rating had gone down to a "5" and the patient's pain score continued to diminish as the wounds healed. Eight weeks after her final wound care visit the patient's skin remained intact and pink without rash. Pruritis was significantly diminished. Six months after her final visit, the skin was still intact!

EPISODE ONE



MAY 22



Lateral wounds of RLE: Moderate serous exudate. Superior malleolus: 0.7 cm x 0.7 cm x 0.3 cm; 50% pink 50% yellow. Distal malleolus: 1.2 cm x 1 cm x 0.5 cm; 20% pink 80% yellow.

Medial wounds of RLE: Moderate serous exudate. Superior: 3.5 cm x 2.7 cm x 0.5 cm; 50% pink 50% yellow slough. Distal: 1.3 cm x 1 cm x 0.3 cm; 50% pink 50% yellow slough.



JULY 3




Lateral wounds of RLE. All areas closed after 6 weeks of treatment.

Medial wounds of RLE: Small amt. serous exudate. Superior: 0.6 cm x 0.3 cm x 0.1 cm; 100% pink; fully granulated.

CONCLUSION

The patient relapsed repeatedly, but this unique treatment (PolyMem Wic and PolyMem Wic Silver cavity filler covered with a 4-layer compression wrap) consistently resulted in improved healing of her wounds. Eventually the combination of using the PolyMem Wic cavity filler plus a 4-layer compression wrap to complete closure and appropriate oral antibiotics resulted in more permanent closure of her wounds.

EPISODE TWO



JULY 31

A month later, new R leg MRSA+ ulcers. Lateral wounds of RLE: 1.3 cm x 1.5 cm x 0.1 cm. Medial & posterior (not shown): 1.5 cm x 1.3 cm x 0.1 cm with numerous other small open areas. Erythemic throughout. Skin is denuded around ulcers. Moderate serous drainage; moderate foul odor.



AUGUST 21

Excellent healing in 3 weeks. Medial wounds: 0.6 cm x 0.5 cm x <0.1 cm; 100% pink. Small amount serous exudate. All other open areas now closed.



AUGUST 28

On oral antibiotics, PolyMem Wic cavity filler & compression. Medial wounds: 0.2 cm x 0.2 cm x 0.1 cm; 100% pink. Small amount serous drainage, no odor, most areas closed.



(Lateral) All wound areas closed. Dismissed.



(Medial) All wound areas closed. Dismissed.

SEPTEMBER 7

POST DISMISSAL

Post-dismissal treatment: Physician placed patient on oral antibiotics 3 times per week for 3 months. Patient applying skin conditioning cream and wearing thigh high compression stockings 20 - 30 mm Hg post-wounds.



MARCH 13

Six months after her final wound care visit, the patient's skin remains intact.

OBJECTIVES

1. Review evidence for the use of PolyMem Wic cavity filler under a 4-layer compression wrap in the treatment of venous hypertension ulcers.
2. Consider the advantages of using PolyMem dressings in terms of passive continuous cleansing of the wound bed, which often eliminates painful and time-consuming wound cleansing during dressing changes.
3. Discuss the benefits of using this unique dressing protocol in terms of quick healing.

This case study was unsponsored. Ferris Mfg. Corp. contributed to the poster presentation.



Ferris Mfg. Corp.

5133 Northeast Parkway | Fort Worth, TX 76106 USA

Toll Free USA: 1.800.POLYMEM (765.9636) | International: +1 630.887.9797

Web site: www.PolyMem.com

BIBLIOGRAPHY:

1. Enoch S, Harding K. Wound bed preparation: The science behind the removal of barriers to healing. *Wounds*. 2003;15(7):213-229.
2. Fowler E, Papen JC. Clinical evaluation of a polymeric membrane dressing in the treatment of dermal ulcers. *Ostomy/Wound Manage*. 1991;35:35-38,40-44.
3. Beitz AJ, Newman A, Kahn AR, Ruggles T, Eikmeier L. A polymeric membrane dressing with antinociceptive properties: analysis with a rodent model of stab wound secondary hyperalgesia. *J Pain*. 2004 Feb;5(1):38-47.

ORIGINAL POSTER PRESENTED AT*:

The WOCN Society's 39th Annual Conference. Poster #1228. June 10 - 13, 2007. Salt Lake City, Utah USA

20th Annual Symposium on Advanced Wound Care (SAWC). Poster #145. April 28 - May 1, 2007. Tampa, FL USA

3rd Congress of the World Union of Wound Healing Societies. Poster #PF405. June 4 - 8, 2008. Toronto, Ontario Canada

* This version has been modified from the original; it reflects PolyMem branding.

PolyMem, PolyMem Silver, PolyMem Wic, Wic, PolyMem Wic Silver, PolyMem Wic Silver Rope, PolyMem Max, Max, PolyMem Max Silver, Shapes, Shapes by PolyMem, The Shape of Healing, The Pink Dressing, SportsWrap, SportsWrapST, More Healing - Less Pain, interlocking circles design, PolyMem For Sports, Not too Loose...Not too Tight...Just Right!, Ferris and FMC Ferris and design are marks owned by or licensed to Ferris. The marks may be registered or pending in the US Patent and Trademark Office and in other countries. Other marks are the property of their respective owners.